

Head Lice



Public Health
Prevent. Promote. Protect.
Hill County Health Department

What are head lice?

Human head lice (*Pediculus humanus capitis*) are parasitic insects that infest the hair on peoples' heads and necks, and rarely on their eyebrows, and eyelashes. Head lice feed on human blood several times a day and live close to the human scalp. Head lice are not known to spread disease.

Who is at risk for getting head lice?

Head lice are found worldwide. In the United States, head lice infestations are seen most often among preschool children attending child care, elementary school children, and household members of infested children, including adults. It is estimated that in the U.S., 6 million to 12 million infestations occur each year among children 3 – 11 years of age.

Head lice cannot hop or fly. They move by crawling and can move very quickly, nearly as fast as you can part someone's hair. Head lice are spread by direct contact with the hair of an infested person. Anyone who has head-to-head contact for any length of time with someone who already has head lice is at risk for infestation. It is uncommon for head lice to be spread by contact with clothing (such as hats, scarves, coats) or other personal items (such as combs, brushes, or towels) used by an infested person. Personal hygiene or cleanliness of the infested person's home or school has nothing to do with getting head lice.

Where are head lice most commonly found?

Head lice and their eggs, called nits, are found almost exclusively on the scalp, particularly around and behind the ears and along the neckline at the back of a person's head. Head lice and nits are rarely found on a person's eyelashes or eyebrows. Head lice hold tightly to hair with hook-like claws at the end of each of their six legs. Head lice nits are cemented firmly to the hair shaft and can be difficult to remove, even after the nymphs hatch and empty casings remain.

What are the signs and symptoms of head lice infestation?

- Tickling feeling of something moving in the hair.
- Itching, after developing sensitivity to the saliva in head lice bites.
- Irritability and difficulty sleeping; head lice are most active at night.

- Sores on the head caused by scratching. These sores can sometimes become infected with bacteria found on the person's skin (head lice, themselves, do not transmit disease).

Six Steps to Getting Rid of Head Lice

1) Examine

Examine everyone in the home for lice and nits.

First, look at a person's hair, close to the scalp around the ears and across the back of the neck. Then carefully check the rest of the head.

Separate the hair strands carefully while looking. Nits are small, white to chocolate brown, oval-shaped eggs that are glued to the hair near the scalp. Nits are more easily seen than live lice.

2) Treat

Treat only those household members who have head lice.

Treat the heads of those you find infested with a lice treatment product. Lice treatment products are available over-the-counter at drugstores and grocery stores. There are some products that can only be prescribed by a medical provider. These products will not kill all of the lice and nits, but they are the best way to get the process started. Carefully follow the directions on the package, or those that your healthcare provider gives you. Using more treatment, reapplying it sooner than recommended, or keeping it on the hair longer than directed will not make it work better or faster and can increase the risk of toxic side effects in the person being treated.

3) Comb

Comb. Comb. Comb.

After treatment, comb the infested person's hair with a lice-removal comb (called a nit comb) to remove lice and nits. Nit combs with long, round, metal teeth tend to be sturdier and more effective. Try to remove every nit and any lice you find. It's OK if you can't get them all on your first try. Just spend time every day until you do get them all.

Combing the hair in the opposite direction of normal brushing will help you find more eggs. Nit picking can also be done by hand, using your fingernails or tweezers. Get rid of nits in the sink, toilet, or garbage. Vacuum around the area where you were combing out nits. Both people should change their clothes afterwards. You can use olive oil to help the comb move through the hair to decrease tangling.

Comb, using the nit comb daily for seven days, or until you no longer find any lice or eggs. This step takes the most time, energy, and patience. **Combing is the most important step!**

4) Clean

Clean your home using normal methods, such as washing clothes, towels, and bedding; vacuuming floors and furniture; and soaking and cleaning combs and brushes in warm, soapy water.

5) Repeat

Most products recommend a second treatment 7-10 days after the first. Follow the directions just as you did for the first treatment. Afterwards, remove any eggs or lice that you might find.

Continue to check heads daily on everyone in your household until one week after the second treatment or one week after the last egg or louse is removed.

6) Notify

Notify everyone that may be affected, for example:

- School
- Church
- Daycare
- Friends
- Family

Head Lice Facts & Tips

- Anyone with clean or dirty hair can get head lice.
- It can be hard work to get rid of them.
- Head-to-head contact is the way head lice are almost spread.
- Lice don't jump or fly, but they are fast crawlers.
- Pets do not carry or spread human lice.
- Head lice do not spread disease.
- Head lice need human blood to live. They'll die within 48 hours if they can't get it.
- It takes head lice eggs (nits) about a week to hatch.
- An itchy scalp is the most common symptom of having head lice and may continue for a while after you have successfully ended an infestation.
- Lice treatment products help kill most of the lice. Even so, you'll still need to physically remove the lice and nits, by combing or picking them out, to be sure they'll no longer be a problem.
- Don't panic or retreat if you see live lice after treatment. Look for them and remove any lice you find until they are all gone. Continue to look until you haven't seen any for 7 days.
- When using NIX, do not apply any type of hair conditioner, hair spray, mousse or gel, or get chlorinated pool water on treated hair for one week.
- The use of mayonnaise, kerosene, Vaseline[®], vinegar, and other home remedies are not proven to

be effective and can be dangerous. Shaving the head is not necessary. If you do not want to use an over-the-counter treatment product, just comb out the nits with a good lice comb.

- Always follow the directions when using treatment products.
- Notify anyone who may be affected.

Head Lice Resources:

American Academy of Pediatrics
National Association of School Nurses
Center for Disease Control